

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/315115884>

Rearing of Adult Green Lacewing, *Chrysoperla carnea* (Stephens) on Different Artificial Diets in the laboratory

Article · January 2016

CITATIONS

6

READS

2,408

11 authors, including:



Aslam Bukero

Sindh Agriculture University

55 PUBLICATIONS 143 CITATIONS

SEE PROFILE



Imtiaz Nizamani

Sindh Agriculture University

39 PUBLICATIONS 140 CITATIONS

SEE PROFILE



Muhammad Ismail Kumbhar

Sindh Agriculture University

26 PUBLICATIONS 44 CITATIONS

SEE PROFILE



Ali Raza Shah

Sindh Agriculture University

25 PUBLICATIONS 75 CITATIONS

SEE PROFILE

Rearing of Adult Green Lacewing, *Chrysoperla carnea* (Stephens) on Different Artificial Diets in the laboratory

Sumera Balouch¹, Aslam Bukero^{1,*}, Imtiaz Ahmed Nizamani¹, Muhammad Ismail Kumbhar², Lubna Bashir Rajput¹, Riaz Ahmed Buriro², Ali Raza Shah³, Naeem Ahmed Qureshi², Javaid Ahmed Sheikh², Shahzad Ali Nahiyoon¹ and Zarnain Rajput¹

¹Faculty of Crop Protection; ²Faculty of Agricultural Social Sciences; ³Faculty of Agricultural Engineering, Sindh Agriculture, University, Tandojam, Pakistan

Abstract: A laboratory study was conducted on rearing of adult Green lacewing, *Chrysoperla carnea* (Stephens) on different artificial diets. Study was conducted under laboratory conditions at Department of Plant Protection, SAU, Tandojam, Sindh, Pakistan during 2013-14 to determined better artificial diet for mass rearing of adult *C. carnea*. Six diets containing water:honey:yeast i.e., D₁ (20ml:2g:2g), D₂ (20ml:4g:2g), D₃ (20ml:6g:2g), water:sugar:yeast D₄ (20ml:2g:2g), D₅ (20ml:4g:2g), D₆ (20 ml:6g:2g) and two diets containing water:gur:yeast i.e., D₇ (20ml:2g:2g), D₈ (20ml:4g:2g), D₉ (20 ml:6g:2g) were tested. The results showed that maximum pre-oviposition period (11.5 ± 3.95 days) recorded, oviposition period (18.0 ± 0.91 days) and post-oviposition period (5.0 ± 0.71 days) was recorded on D₄, D₉ and D₁, respectively. Minimum pre-oviposition period (7.25 ± 0.25 days), oviposition period (5.0 ± 1.73 days) and post-oviposition period (2.5 ± 0.87 days) was recorded on D₈, D₄, and D₇, respectively. The highest fecundity (342.5 ± 89.55 eggs) was recorded on D₉, whereas the lowest fecundity (42.5 ± 5.48 eggs) was recorded on D₄. The eggs laid by female fed on D₄ took lesser time (1.75 ± 0.66 days) to hatch. Maximum larval and pupal longevity was recorded in larvae of adults fed on D₉. The adults lived longer for 30.0 ± 0.91 days on D₉. The maximum egg hatching (55 %), larval-pupal transformation (45 %) and adults emergence (30 %) was recorded on D₉. It is concluded that D₉ was found the best one for mass rearing of adult *C. carnea*.

Keywords: *C. carnea*, Development time, Biology, Survival, Artificial diets.

INTRODUCTION

The horticultural and field crops are attacked by arthropod pests, which deteriorate the quality and quantity of yields and force. Farmers to use agrochemicals. Green lacewing, *Chrysoperla carnea* Stephens (Neuroptera: Chrysopidae) is mass-reared and used as biocontrol agent against soft bodied insect pest [1]. Its larvae feeds on insect pest of field as well as fruit crops. This predator was mass reared and augmented as biological control agents against several arthropods in American as well as European countries. It is commercially available and released as individual stages (eggs, larvae or pupae) or combined stages, according to the demand of the consumers. Adult stage of *C. carnea* mass-reared on artificial foods i.e. yeast, sugar products and water boosted their activeness at large scale [2]. Adults of *C. carnea* were successfully mass-reared when fed on artificial diets including yeast, pollen and honey [3]. The diets contained artificial ingredients significantly influenced oviposition and post-oviposition periods and fecundity of females [4,5]. However, when *C. carnea* larvae were reared on nymphs of *Shizaphis graminum*, *Bemisia tabaci* as well as artificial diets, significant impact was determined on fecundity, larval duration, hatchability and sex ratio

(female/male). During the scarcity of natural foods the artificial diet is the best source for mass rearing of *C. carnea* [6]. The present research work was conducted on different artificial food supplements for mass rearing of *C. carnea* adults in the laboratory. The findings of present result will be beneficial for mass production and augmentation of adult *C. carnea* to regulate the insect pest populations on horticultural and field crops.

MATERIALS AND METHODS

The research experiment was conducted on "Rearing of Green lace wings, *C. carnea* on different artificial diets" during 2013-14 in the laboratory of Plant Protection Department, Sindh Agriculture University, Tandojam. The experiment was laid down as Randomized Complete Design (CRD) with five replications each having three pairs of adult *C. carnea*. These adults were confined in the glass chimney (6 cm Ø × 8 cm Ø) and placed in the petri dish (9.0 cm Ø). A small glass vial was placed inside the chimney holding cotton soaked in distilled water to maintain moisture. The upper open end of glass chimney was covered with black muslin cloth and was tightened with rubber band. The different adult diets was provided inside the glass chimney with the help of dropper on glass strips, each strip being drilled at three points to make pits for holding drops of diet. The diets were provided on daily basis to the adults. The eggs was laid by females on the walls of chimney and muslin cloth was removed

*Address correspondence to this author at the Department of Entomology, Sindh Agriculture University, Tando Jam, Pakistan; Tel: 923003085339; E-mail: bukero786@gmail.com

with sharp razor and ten egg per test tube (10 × 1 cm Ø) was placed with the help of camel hair brush. The test tubes were covered with cotton swab. After hatching, the newly emerged larvae were reared on frozen eggs of *Sitotroga cerealella* (Oliver) (Lepidoptera: Gelechiidae). Nine different diets were tested for *C. carnea* adults. The ratio of different ingredients was measured on electric weight balance according to the composition of diets. Six diets containing water:honey:yeast i.e., D₁ (20ml:2g:2g), D₂ (20ml:4g:2g), D₃ (20 ml:6g:2g), water:sugar:yeast D₄ (20ml:2g:2g), D₅ (20ml:4g:2g), D₆ (20 ml:6g:2g) and three diets containing water:gur:yeast i.e., D₇ (20ml:2g:2g), D₈ (20ml:4g:2g), D₉ (20 ml:6g:2g) were tested. The experiment was conducted at 26±2°C and 60±5 % RH as reported by [7]. Daily observation were carried out to know the fecundity, the duration in the days of pre-oviposition, oviposition and post-oviposition periods, the egg hatching percentage, the larval development duration, pupal duration, survival and mortality on different artificial diets. The collected data was subjected for statistical analysis and statistical differences existed between data sets ($P < 0.05$), Fisher's Least Significant Differences (LSD) was used to separate the differing means [8].

RESULTS

The results given in Table 1 indicated that the maximum pre-oviposition period was recorded as (11.5 ± 3.95) days on diet D₄ followed by 10.75 ± 1.38, 10.0 ± 0.41, 10.0 ± 2.0, 9.0 ± 0.41, 8.0 ± 1.0, 8.0 ± 0.41, 7.25 ± 1.35 and 7.25 ± 0.25 on diets D₅, D₆ and D₇, D₂, D₃, D₁ and D₈ days, respectively. The maximum oviposition

period was observed as (18.0 ± 0.91) days on diet D₉ followed by 16.5 ± 2.03, 15.0 ± 1.94, 12.5 ± 1.44, 9.5 ± 3.12 and 5.0 ± 1.73 days on diets D₃, D₇ and D₈, D₂, D₁, D₅ and D₆ and D₄, respectively. The highest fecundity was recorded 342.5 ± 89.55 eggs when adults fed with diet D₉ followed by 163.25 ± 49.40, 162.5 ± 6.37, 116.25 ± 5.39, 110.75 ± 25.81, 74.0 ± 12.42, 63.5 ± 12.59, 53.75 ± 9.28 and 42.5 ± 5.48 on diets D₈, D₇, D₇, D₃, D₁, D₆, D₅ and D₄, respectively. The maximum post-oviposition period was observed as 5.0 ± 0.71 days on diet D₁ followed by 4.5 ± 0.65, 4.0 ± 0.41, 3.75 ± 0.48, 3.5 ± 0.29, 3.0 ± 0.41 and 2.5 ± 0.87 on diets D₆, D₉, D₈, D₃, D₂, D₄, D₅ and D₇, respectively. The highest adult survival (30.0 ± 0.91) days as was recorded in predators reared on the diet D₉ followed by 29.0 ± 2.69, 28.0 ± 2.65, 27.0 ± 1.29, 26.0 ± 2.55, 25.0 ± 2.80, 24.0 ± 0.41, 22.5 ± 1.50 and 19.0 ± 3.0 on diets D₇, D₃, D₈, D₂, D₁, D₆, D₅ and D₄, respectively.

The results in Table 2 depicted the maximum incubation period was observed as (6.25 ± 1.25) days of the egg laid by adult when fed on the diet D₃ followed by 5.0 ± 1.12, 4.5 ± 1.06, 4.0 ± 1.0, 3.75 ± 0.97, 3.0 ± 0.87 and 1.75 ± 0.66 on diets D₇, D₉, D₅, D₆, D₁ as well as D₂ and D₄, respectively. The data further revealed that the maximum survivorship of 1st instar larva was observed as (3.25 ± 0.90) days on the diets D₃ as well as D₃ followed by 3.0 ± 0.87 and 2.75 ± 0.83 and 1.75 ± 0.66 days on diets D₉ as well as D₁ and D₅ as well as D₆ and D₄, respectively. Similarly the maximum survivor (3.25 ± 0.90) days of 2nd instar larva was recorded on the diets D₉ as well as D₇ followed by 3.0 ± 0.87 and 2.75 ± 0.83, 2.5 ± 0.79, 2.25 ± 0.75 and 1.75 ± 0.66 days on diets D₈ as well as D₆, D₅, D₁ as well as D₃ and D₄, respectively. The 3rd instar larva lived

Table 1: Response of *Chrysoperla carnea* Adults on Different Artificial Diets Under Laboratory Conditions

Diets	Biological Parameters				
	Pre-oviposition (Days)	Oviposition (Days)	Fecundity (No. eggs/female)	Post-oviposition (Days)	Survival (Days)
D ₁	7.25 ± 1.35 e	12.5 ± 1.44 d	74.0 ± 31.42 e	5.0 ± 0.71 a	25.0 ± 2.80 c
D ₂	8.0 ± 1.00 d	15.0 ± 1.94 bc	110.75 ± 25.81 d	3.0 ± 0.41 c	26.0 ± 2.55 c
D ₃	8.0 ± 0.41 d	16.5 ± 2.03 b	116.25 ± 5.39 c	3.5 ± 0.29 bc	28.0 ± 2.65 ab
D ₄	11.5 ± 3.95 a	5.0 ± 1.73 f	42.5 ± 31.48 h	2.5 ± 0.87 c	19.0 ± 3.00 e
D ₅	10.75 ± 1.38 a	9.5 ± 3.12 e	53.75 ± 23.28 g	2.5 ± 0.87 c	22.5 ± 1.50 d
D ₆	10.0 ± 0.41 b	9.5 ± 0.65 e	63.5 ± 12.59 f	4.5 ± 0.65 b	24.0 ± 0.41 c
D ₇	10.0 ± 2.00 b	16.5 ± 2.03 b	162.5 ± 6.37 b	2.5 ± 0.29 c	29.0 ± 2.69 a
D ₈	7.25 ± 0.25 e	16.5 ± 1.55 b	163.25 ± 49.40 b	3.75 ± 0.48 b	27 ± 1.29 ab
D ₉	9.0 ± 0.41 c	18.0 ± 0.91 a	342.5 ± 89.55 a	4.0 ± 0.41 b	30.0 ± 0.91 a

Different letters within a row indicate significant difference (Fisher's Protected LSD test: $P < 0.05$).

Table 2: Mean Developmental Duration in Days of Immature Life Stages of *Chrysoperla carnea* from Females Fed on Different Artificial Diets under Laboratory Conditions

Life Stages	Artificial diets								
	D1	D2	D3	D4	D5	D6	D7	D8	D9
Egg	3.0 ± 0.87 ijkl	3.0 ± 0.87 ijkl	6.25 ± 1.25 cd	1.75 ± 0.66 l	4.0 ± 1.00 ghi	3.75 ± 0.97 ghij	5.0 ± 1.12 defg	4.0 ± 1.00 ghi	4.5 ± 1.06 fgh
1 st	2.75 ± 0.83 ijkl	3.0 ± 0.87 ijkl	3.25 ± 0.90 hijk	1.75 ± 0.66 l	2.75 ± 0.83 ijkl	2.75 ± 0.83ijkl	3.25 ± 0.90 hijk	2.75 ± 0.83 ijkl	3.0 ± 0.87 ijkl
2 nd	2.5 ± 0.79 jkl	2.25 ± 0.75 kl	2.5 ± 0.79 jkl	1.75 ± 0.66 l	2.75 ± 0.83 ijkl	3.0 ± 0.87ijkl	3.25 ± 0.90 hijk	3.0 ± 0.87 ijkl	3.25 ± 0.90 hijk
3 rd	2.5 ± 0.79 jkl	2.25 ± 0.75 kl	2.5 ± 0.79 jkl	1.75 ± 0.66 l	2.75 ± 0.83 ijkl	2.5 ± 0.79 jkl	3.0 ± 0.87 ijkl	3.0 ± 0.87 ijkl	2.75 ± 0.83 ijkl
Pupa	6.0 ± 1.22 cde	5.5 ± 1.17 def	4.75 ± 1.09 efg	2.5 ± 0.79 jkl	7.75 ± 1.39 ab	9.0 ± 1.50 a	8.75 ± 1.48 a	7.0 ± 1.32 bc	9.0 ± 1.50 a

Different letters within a row indicate significant difference (Fisher's Protected LSD test: $P < 0.05$).

Table 3: Egg hatching, larval-pupal transformation and adult emergence % of *Chrysoperla carnea* individuals from females fed on different artificial diets under laboratory conditions

Life stages	Artificial diets								
	D ₁	D ₂	D ₃	D ₄	D ₅	D ₆	D ₇	D ₈	D ₉
Hatch %	42.5	42.5	27.5	20.0	35.0	37.5	40.0	55.0	55.0
Pupa	17.5	20.0	15.0	12.5	30.0	30.0	32.5	45.0	40.0
Adult	17.5	12.5	10.0	7.5	17.5	22.5	20.0	27.5	30.0

for more days (3.0 ± 0.87) on the diets D₇ as well as D₈ as compared to 2.75 ± 0.83 , 2.5 ± 0.79 , 2.25 ± 0.75 and 1.75 ± 0.66 days on diets D₅, D₉, D₁, D₃, D₂ and D₄, respectively. The results further depicted that the maximum survivorship was seen in the pupal stage as (9.0 ± 1.50) days on the diets D₆ as well as D₉ followed by 8.75 ± 1.48 , 7.75 ± 1.39 , 7.0 ± 1.32 , 6.0 ± 1.22 , 5.5 ± 1.17 , 4.75 ± 1.09 and 2.5 ± 0.79 days on diets D₇, D₅, D₈, D₁, D₅, D₃ and D₄, respectively. The result further revealed that the percentage of egg hatching, larval-pupal transformation and adult emergence was seen on different artificial diets. The data in Table 3 showed that the maximum hatching percentage was recorded 55 % on the diets D₈ as well as D₉ followed by 42.5 %, 40.0 %, 37.5 %, 35.0 %, 27.5 % and 20.0 % on diets D₁ as well as D₂, D₇, D₆, D₃ and D₄, respectively. It was further observed that the maximum pupal percentage was recorded 45 % on diet D₈ followed by 40 %, 32.5 %, 30.0 %, 20.0 %, 17.5 %, 15.0 % and 12.5 % on diets D₉, D₇, D₅ as well as D₆, D₂, D₁, D₃, and D₄, respectively. The highest adult emergence was recorded as (30 %) on diet D₉ followed by 27.5 %, 22.5 %, 20.0 %, 17.5 %, 12.5 %, 10.0 % and 7.5 % on diets D₈, D₆, D₁ as well as

D₅, D₂, D₃ and D₄, respectively. The analysis of variance showed that there was significant difference on diets and life stages ($P < 0.05$).

DISCUSSION

The findings of present result have the conformity with McEwen and Kidd [2] who reported that *C. carnea* significantly lived longer on diet containing sugar only as compared to sugar and yeast ($P < 0.05$). The highest fecundity was obtained the diet containing yeast and when artificial diets containing yeast, sugar and water (4:7:10) ratios, egg production was significantly reduced in small numbers. Our findings have partial agreements with those of Nawaz *et al.* [9] who reported that adults fed on diet contained vitamin E were found most suitable for adult *C. carnea* rearing. Both male and female adults lived longer. The same results were also achieved when adults reared on Ber honey followed by honey obtained from almond, orange and apple trees. Our findings have the partial conformity with those of Syed *et al.* [10] determined the nine diets provided to immature stages of *C. carnea*, including six

diets were containing vitamins and minerals and rest of three were natural. The result revealed that shortest total larval period 13.9 days with maximum cocoon weight (0.828 g), and fecundity were determined 717 eggs reared on Diet 9 (*S. cerealella* eggs). The highest larval duration 27.65 days were obtained on Diet 4 and maximum mortality 65% were seen on Diet 3. Sarailoo and Lakzaei [11] determined the impact of six artificial diets contained [a mixture of 30% concentrations of glucose, fructose and sucrose (1: 1: 1), glucose, fructose, sucrose plus extract of *S. cerealella* eggs (1: 1), glucose, fructose, sucrose plus extract of *A. kuehniella* eggs (1: 1), a mixture of honey, yeast and distilled water (1: 1: 1), honey, yeast plus extract of *S. cerealella* eggs (1: 1: 1) and honey, yeast plus extract of *A. kuehniella* eggs (1: 1: 1)]. The findings of this study showed that the mixture of honey, yeast and extract of *A. kuehniella* eggs (1: 1: 1), were found best results on biological aspects of the predator as compare to other diets.

CONCLUSION

The maximum pre-oviposition, oviposition and post-oviposition period was observed on diet D₄, D₉ and D₁, respectively. The highest fecundity was obtained on diet D₉, whereas, lowest on diet D₄. The minimum egg incubation period, larval and pupal development was recorded on diet D₄, whereas, maximum was recorded on diet D₉. The highest survivor of adults was seen on diet D₉, however, the lowest survivor was obtained on diet D₄. The maximum egg hatching, pupal and adult emergence percentage was observed on diet D₉, whereas, the minimum was recorded on diet D₄.

REFERENCES

- [1] Ulhaq MM, Sattar A, Salihah Z, Farid A, Usman A, Khattak SUK. Effect of different artificial diets on the biology of adult green lacewing (*Chrysoperla carnea* Stephens.) Songklanakarin Jr Sci Technol 2006; 28(1): 1-8.
- [2] McEwen KP, Kidd NAC. The Effects Of Different Components Of An Artificial Food On Adult Green Lacewing (*Chrysoperla carnea*) Fecundity And Longevity. Entomol Experimentalis et Applicata 1995; 77(3): 343-346. <http://dx.doi.org/10.1111/j.1570-7458.1995.tb02332.x>
- [3] Ribeiro LJ, Freitas SD. Influence of food on *Chrysoperla externa* (Hagen) reproductive potential. Revista-de-Agricultura- Piracicaba 2000; 75(2): 187-196.
- [4] Milevoj L. Rearing of the common Green lacewing (*Chrysoperla carnea* Steph.) In the laboratory. Zbornik. Biotehniške. Fakultete. Univerze v. Ljubljani. Kmetijstvo. 1999; 73: 65-70.
- [5] Adane T, Gautam RD. Biology and feeding potential of green lacewing, *Chrysoperla carnea* on rice moth. Indian Jr Entomol 2002; 64(4): 457-464.
- [6] Jokar M, Zarabi M. Investigation Effect Three Diets On Life Table Parameters *Chrysoperla Carnea* (Steph.) (Neuroptera: Chrysopidae) Under laboratory Conditions. Egypt Acad Jr Biolog Sci 2012; 5(1): 107-114.
- [7] Ashfaq AM, Nasreem B, Cheema GM. Advances In Mass Rearing Of *Chrysoperla carnea* (Stephen) (Neuroptera: Chrysopidae). South Pacific Study 2004; 24(2): 47-53.
- [8] Zar JH. Biostatistical Analysis. Prentice-Hall, Upper Saddle River, N. J, USA 1996; p. 662.
- [9] Nawaz M, Ashfaq M, Ali A. Studies on improvement of artificial diet and its effect on biological characters of *Chrysoperla carnea* (Stephens). Pak Entomol 2008; 30(1): 73-76.
- [10] Syed AN, Ashfaq M, Ahmad S. Comparative Effect of various diets on development of *Chrysoperla carnea* (Neuroptera: Chrysopidae). Int Jr Agric Biol 2008; 10(6): 728-30.
- [11] Sarailoo MH, Lakzaei M. Effect of Different Diets On Some Biological Parameters Of *Chrysoperla carnea* (Neuroptera: Chrysopidae). Jr Crop Prot 2014; 3(4): 479-486.

Received on 24-02-2016

Accepted on 12-04-2016

Published on 20-07-2016

<http://dx.doi.org/10.6000/1927-5129.2016.12.44>

© 2016 Sumera et al.; Licensee Lifescience Global.

This is an open access article licensed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0/>) which permits unrestricted, non-commercial use, distribution and reproduction in any medium, provided the work is properly cited.